


INTIMATE PARTNER VIOLENCE AND CONTRACEPTIVE USE DISCONTINUATION IN INDIA

Intimate Partner Violence (IPV) in India

About 33% of married women ages 15-49 in India have experienced IPV in their lifetime — spousal physical, emotional, or sexual violence.

(IIPS and ICF 2017)



Contraceptive Use Discontinuation

50% of Indian women who recently discontinued contraceptive use for 1+ month still had a need for it.

! IPV can compromise women's reproductive self-efficacy.

! The relationship between IPV and contraceptive discontinuation is highly variable and poorly understood.

Indian women who relied on **IUDs** for contraception and experienced

PHYSICAL PARTNER VIOLENCE

had more than **3X RISK** of discontinuing contraceptive use while still in need.



Indian women who relied on **MALE CONDOMS** for contraception and experienced

EMOTIONAL VIOLENCE

had more than **4X RISK** of discontinuing contraceptive use while still in need.



Understanding women's experiences of violence is critical to supporting the achievement of their reproductive goals.

Details

Infographic is derived from research presented in: Upadhyay A, Kumar K, James KS, McDougal L, Raj A, Singh A. (2022). Association between Intimate Partner Violence and Contraceptive Use Discontinuation in India. Studies in Family Planning.

Contact

iipsgehproject@gmail.com

THE GENDER PROJECT
THE GENDER EQUITY AND DEMOGRAPHIC RESEARCH PROJECT



GEH
GENDER EQUITY AND DEMOGRAPHIC RESEARCH PROJECT